



Spectrum Journal of Innovation, Reforms and Development

Volume 38, April 2025

ISSN (E): 2751-1731

WEBSITE: WWW.SJIRD.JOURNALSARK.ORG

DEVELOPING NUMBER AND DATE OF SORGHUM IN THE CONDITION OF SYRDARYA PROVINCE

Ch. V. Toshpulatov 1

B. B. Tukhtashev 1

B. T. Mavlonov 2

1Tashkent State Agrarian University, University str. 2,
Tashkent Province, 100140, Uzbekistan

2Samarkand Veterinary Medicine Institute,
Ulugbek Street, 77, Samarkand, 130100, Uzbekistan

Abstract

It is no secret that the rapid land reclamation, i.e., their salinization and depletion of irrigation water reserves, will turn Uzbekistan into the most difficult region of the arid region in the coming years. Now it is appropriate to conduct large-scale practical and scientific research on improving the land reclamation state of soils and saving irrigation water reserves in the republic's agriculture (4, 5, 10).

During the visit of the President of the Republic of Uzbekistan to the Syrdarya region in May 2017 and April 2018, the "Bek Cluster" enterprise was established, where it is planned to grow 80 thousand tons of fodder crops. In order to achieve the specific task set out above, the cultivation of fodder crops in the conditions of the Syrdarya province is an urgent issue (1).

Introduction

METHODS AND MATERIALS

In field experiments in agricultural crops, the author Dospekhov B.A. "Methodology of conducting field experiments" developed by M. Kolos. 1979y. and Nurmatov Sh. "Field Experiment Methodology" developed by and others used (2, 3). Based on the natural climatic conditions of the "Oq Oltin" district of the Syrdarya region, medium salinity land is selected for the field experiment.

The field experiment was carried out in the following options:

In options 1-3, the soil salt was not washed. Sorghum were watered at 70-80-75%; 70-80-70%; 70-70-70%;

In options 4-6, soil salinity was washed in November. Sorghum were irrigated at 70-80-75%; 70-80-70%; 70-70-70% compared to LFMC;

In options 7-9, soil salinity was washed away in February. Sorghum were irrigated at 70-80-75%; 70-80-70%; 70-70-70% compared to LFMC. When sorghum is planted in rows, 12 kg/ha is the norm. The row spacing is 60 cm. The plant spacing is 10 cm. Planting gives good results.



The agrochemical and water properties of the soil are studied in the experimental field. In order to determine the agrochemical parameters of the soil of the experimental field, mixed soil samples were taken from 0-30 and 30-50 cm soil layers by envelope method from 5 points of the field. The total amount of humus and humus in these samples I.M. Tyurin; nitrogen and phosphorus I.M. Maltseva, L.N.Gritsenko; in a nitrate nitrogen-ionometric instrument; mobile phosphorus is determined by the methods of B.P. Machigin and exchangeable potassium by P.V.Protasov.

Samples were taken in the 0th and 30th 50 cm layers of the plowed and under-plowed soil to determine the amount of NPK, general and mobile forms, humus, humus and sent to the laboratory for analysis (2, 3, 4) 30 cm. The volume weight of the soil is determined according to the irrigation procedures in each layer at depths of 0-10-50 cm. The water permeability of the soil was determined using special cylinders in the spring and after harvesting.

RESULTS AND DISCUSSION

In the experiment, sorghum was irrigated 3 times out of 5 during the growing season depending on soil moisture deficiency. The most important indicator is determining the period of irrigation of sorghum. Before the next irrigation, the actual soil moisture and the limiting field moisture capacity (LFMC) were very useful in determining the correct timing of irrigation. Each successive irrigation was determined based on the actual soil moisture and the limit field moisture capacity (LFMC) indicator.

In 2019, in the control variant of the experiment, the 1st irrigation fell on June 4th, the 2nd irrigation fell on June 25th, the 3rd irrigation fell on July 7th, and the 4th irrigation fell on July 25th, and finally the 5th irrigation fell on August 10th. The period between successive irrigations was 21-22 days during the pre-irrigation soil moisture content of 70% compared to LFMC in option 1, and 12-13 days during the pre-irrigation soil moisture content of 80% compared to LFMC. In this variant, the soil moisture during the growing season was 70-80-75% in comparison with LFMC.

In the second option, the 1st irrigation was on June 4th, the 2nd irrigation was on June 26th, the 3rd irrigation was on July 7th, and finally the 4th irrigation was on August 1st. Based on that, the 1st irrigation was given on 22 days, the 2nd irrigation on 12-13 days, the 3rd irrigation on 18-19 days, and the 4th irrigation after 19-20 days. In this option, soil moisture was maintained at 70-80-70% during the growing season compared to LFMC.

In the 3rd option, where the soil salt was not washed, the 1st irrigation of sorghum was on June 4, the 2nd irrigation was on July 26, and finally the 3rd irrigation was on July 22. The period between irrigations was 25-26 days. In this option, soil moisture was maintained at 70-70-70% during the vegetation period compared to LFMC.

In variant 4, which was washed with soil salinity in November, the 2nd irrigation was carried out on June 24, the 3rd irrigation on July 9, the fourth irrigation on July 28 and the 5th irrigation on August 12. The period between successive irrigations was 19-20 days in the period when the pre-irrigation soil moisture was 70% compared to LFMC in option 4, and 14-15 days in the period when the pre-irrigation soil moisture was 80% compared to LFMC was kept at 70-80-75%.

In option 5, the 2nd irrigation was carried out on June 25, the 3rd irrigation on July 10, and the fourth irrigation on August 2. The period between successive irrigations was 19-22 days when the



soil moisture was 70% relative to LFMC during the growing season, and 14-15 days when the soil moisture was 80% relative to the LFMC during the growing season. In this option, soil moisture was maintained at 70-80-70% during the growing season compared to LFMC.

In the 6th option, the 1st irrigation of sorghum fell on June 5, the remaining irrigations, including the 2nd irrigation, were carried out on July 27, and finally, the 3rd irrigation was carried out on July 21. The period between the next irrigations was 24-25 days in the fourth option. In this option, the soil moisture was kept at 70-70-70% compared to LFMC during the growing season.

In option 7, which was washed with soil salinity in February, the 2nd irrigation was carried out on June 25, the 3rd irrigation on July 10, the fourth irrigation on July 29 and the 5th irrigation on August 14. The period between successive irrigations in the 7th option was 19-20 days in the period when the pre-irrigation soil moisture was 70% compared to LFMC, and 14-15 days in the period when the pre-irrigation soil moisture was 80% compared to LFMC. In this variant, soil moisture was kept at 70-80-75% during the growing season compared to LFMC.

In option 8, the 2nd irrigation was carried out on June 27, the 3rd irrigation on July 11, and the fourth irrigation on July 3. The period between successive irrigations was 20-21 days in the period when the pre-irrigation soil moisture was 70% compared to LFMC in option 8, and 14-15 days in the period when the pre-irrigation soil moisture was 80% compared to LFMC. In this option, the soil moisture during the vegetation compared to LFMC was kept at 70-80-70%.

In the 9th option, the 2nd irrigation was carried out on June 28, and the 3rd irrigation was carried out on July 23. The period between the next irrigations was 24-26 days in the seventh option. In this option, the soil moisture was kept at 70-70-70% during the vegetation period compared to LFMC.

In the control variant, due to the fact that the soil salt was not washed, the process of coming to the next water of sorghum was accelerated, because during the growing season of the plant, there was not enough moisture in the soil for the growth and development of the plant (6, 7).

Table 2 The dates, number and period between irrigations of sorghum. Data for 2019.

Option No	1 st irrigation		2 nd irrigation		3 th irrigation		4 th irrigation		5 th irrigation	
	irrigation period	day	irrigation period	day	irrigation period	day	irrigation period	day	irrigation period	day
1	2	3	4	5	6	7	8	9	10	11
1.	4.06	21-22	25.06	12-13	7.07	18-19	25.07.	16-17	10.08	
2.	4.06	21-22	26.06	12-13	7.07	18-19	1.08		-	
3.	4.06	21-22	26.06	25-26	19.07	-	-		-	
4.	5.06	19-20	24.06	14-15	9.07	21-22	27.07	16-17	12.08	
5.	5.06	19-20	25.06	14-15	10.07	21-22	2.08			
6.	5.06	21-22	27.06	24-25	21.07					
7.	6.06	19-20	25.06	14-15	10.07	18-19	29.07	15-16	14.08	
8.	6.06	20-21	27.06	13-14	11.07	21-22	3.08			
9.	6.06	21-22	28.06	24-25	23.07					

**Table 3 The date, number and period between irrigations of sorghum. Data for 2020.**

Option No	1 st irrigation		2 nd irrigation		3 th irrigation		4 th irrigation		5 th irrigation	
	irrigation period	day	irrigation period	day	irrigation period	day	irrigation period	day	irrigation period	day
1	2	3	4	5	6	7	8	9	10	11
1.	9.06	19-20	28.06	13-14	12.07	14-15	27.07	15-16	12.08	
2.	9.06	19-20	29.06.	14-15	14.07	15-16	29.08		-	
3.	9.06	22-23	2.07	23-24	25.07		-		-	
4.	10.06	18-19	29.06	14-15	14.07	13-14	28.07.	14-15	14.08	
5.	10.06	19-20	30.06	14-15	15.07	14-15	30.07			
6.	10.06	22-23	3.07	25-26	27.07					
7.	11.06	19-20	2.07.	12-13	15.07	14-15	30.07.	14-15	14.08	
8.	11.06	18-19	29.06	14-15	14.07	14-15	1.08			
9.	11.06	22-23	3.07	25-26	28.07					

Table 4 The duration, number and period between irrigations of sorghum. Data for 2021

Option No	1 st irrigation		2 nd irrigation		3 th irrigation		4 th irrigation		5 th irrigation	
	irrigation period	day	irrigation period	day	irrigation period	day	irrigation period	day	irrigation period	day
1	2	3	4	5	6	7	8	9	10	11
1.	7.06	18-19	26.06.	13-14	8.07	14-15	23.07	16-17	10.08	
2.	7.06	18-19	25.06	14-15	9.07	18-19	28.07		-	
3.	7.06	23-24	1.07	23-24	25.07		-		-	
4.	8.06	17-18	25.06	13-14	9.07	14-15	24.07	16-17	12.08	
5.	8.06	17-18	25.06	14-15	10.07	18-19	29.07			
6.	8.06	22-23	2.07	22-23	25.07					
7.	9.06	16-17	25.06.	14-15	9.07	14-15	24.07	16-17	12.08	
8.	9.06	16-17	26.06	15-16	12.07	19-20	2.08			
9.	9.06	22-23	2.07	23-24	26.07				-	

In experimental options (options 4, 5, 6) where the soil salt was washed in November, the next irrigation period was delayed by 2-3, and in some cases, 4-5 days. The main reason for this is that the soil is provided with a sufficient amount of moisture reserves in the process of washing the soil salt, and the loss of this moisture caused a slight delay in the irrigation period. According to our analysis, in these options (options 4, 5, 6), the soil capillaries were restored due to the washing away of harmful salts from the soil, and in early spring, the moisture in the lower layer was accelerated up through the capillary pores, due to which the moisture reserve in the soil reached the next irrigation.

In the experiment, it was noted that the next irrigation was slightly delayed due to the washing of soil salt in February. The main reason for this is that the moisture in the soil was preserved for a long time during the washing of the soil salt in February, and the loss of this moisture caused a slight delay in the irrigation period (9).

In the experiment, the sorghum used enough of the soil moisture reserves from the time it sprouted until flowering. We can also know this from the water consumption indicators of the sorghum. In the experiment, it was reported that sorghum satisfied its water demand by 30-33% from rain and



reserve water (7,8) Only one It should be noted that in 2019-2020, when the soil moisture was 70% relative to LFMC, the period between irrigation was 21-22 days, and when it was 80% relative to LFMC, the period between sorghum irrigation was 14-15 days, in 2021, the period between sorghum irrigations was 1 It was reduced by 2 days or the plant came to water early. This year, precipitation was 30-40% less than in 2019-2020.

In these years, when the water shortage is increasing, keeping the soil moisture at the level of 75-80% LFMC during the growing season raises the level of seepage water, makes it possible to re-salinize the soil, and worsens its amelioration condition. In order for sorghum to properly use soil moisture during its growth and development, it is advisable to schedule irrigation from 14-15 days to 19-20 days based on the plant's development phases.

REFERENCES

1. Decree No. PF-4947 of the President of the Republic of Uzbekistan dated February 7, 2017 "On the strategy of actions for the further development of the Republic of Uzbekistan". Collection of legal documents of the Republic of Uzbekistan, 2017. No.6, Article 70.
2. Field experiments transfer styles. Tashkent: UzPITI, 2007. P.146
3. Vavilov P.P., Rasteniyevodstvo, Moscow "Kolos"-1979. str 514
4. Yormatova D., Ubaidullaev Sh., Halimov I. Tokhtashev B., Khushvaktova H. Cultivation of field crops. Tashkent Science Publishing House-2004.
5. Mirzaev O.F., Khudoyberdiyev T.S., Cultivation of fodder. Andijan publishing house, Tashkent-2003.
6. Norkulov U. Effective use of water in salt washing (recommendation) Tashkent-2018.
7. Tokhtashev B.B., Yormatova D. , Abduazizov B. Agricultural activities of the winter season on farms. Farmer's journal., Tashkent 2014.
8. Norkulov U., Tukhtashev B., Eshunkulov J. Change of Mechanical Composition of Soils after Flood of Sardoba Water Reservoir International Journal of Innovative Analyzes and Emerging Technology e-ISSN: 2792-4025 | [http:// openaccessjournals. EU](http://openaccessjournals.eu) | Volume: 2 Issue: 2
9. Tukhtashev B., Toshpulatov Ch, Shadmonov, M Khudaybergenov N. Role and productivity of white sorghum in the reclamation of saline soils of Uzbekistan., E3S Web of Conferences 258,04053 (2021) UESF-2021, page 7
10. Toshpulatov Ch, Tukhtashev B, Charshanbiev U and Mavlonov B. Effects of soil salt-leaching terms on growth, development and yield of sorghum in Uzbekistan. IOP Conference Series: Earth and Environmental Science. To cite this article: Toshpulatov et al. 2022 IOP Conf. Ser.: Earth Environ. Sci.